

Fusion systems on metacyclic 2-groups

Benjamin Sambale
 Mathematisches Institut
 Friedrich-Schiller-Universität
 07743 Jena
 Germany
 benjamin.sambale@uni-jena.de

August 6, 2009

Abstract

Let P be a finite metacyclic 2-group and $\mathcal{F}$ a fusion system on P . We prove that $\mathcal{F}$ is nilpotent unless P has maximal class or P is homocyclic, i. e. P is a direct product of two isomorphic cyclic groups.

1 Introduction

The structure of a finite group G is often controlled by the structure of its Sylow p -subgroups and the way they are embedded into G . In this paper, we give an example of such a control in the case $p = 2$. More precisely, we show that G is 2-nilpotent if the Sylow 2-subgroups of G are metacyclic, but neither of maximal class nor homocyclic. Here, a homocyclic group means a direct product of two isomorphic cyclic groups. Moreover, the 2-groups of maximal class are precisely the dihedral groups, the semidihedral groups, and the quaternion groups. Of course, there are many other metacyclic 2-groups (see for example [3]). In this sense, most of the metacyclic 2-groups satisfy our assumption. In the easiest case where the Sylow 2-subgroups of G are cyclic, the result is a well-known theorem by Burnside. Another case, in which the result is known, is due to Wong. He showed that G is 2-nilpotent if the Sylow 2-subgroups of G are the sometimes called modular groups (see Satz IV.3.5 in [2]). These are the nonabelian 2-groups with cyclic subgroups of index 2, which do not have maximal class. For similar results see [7].

For the proof, we use the notion of fusion systems, which provide a great generalization of the situation described above. In particular, our result applies also to 2-blocks of finite groups with metacyclic defect groups. We note that David Craven and Adam Glesser also found the main result of this paper, but they did not published it yet. Furthermore, Radu Stancu has studied fusion system on metacyclic p -groups for odd primes p (see [6]).

2 Metacyclic 2-groups

First, we collect some elementary results about metacyclic 2-groups. We will freely use the fact that subgroups and quotients of metacyclic groups are again metacyclic.

Lemma 1. *If P is a metacyclic 2-group, then the automorphism group $\text{Aut}(P)$ is also a 2-group unless P is homocyclic or a quaternion group of order 8.*

Proof. See Lemma 1 in [5]. □

It is easy to show that the converse of Lemma 1 also holds. In the following, we denote the cyclic group of order $n \in \mathbb{N}$ by C_n . Moreover, we set $C_n^2 := C_n \times C_n$. For a finite group G , we denote by $\Phi(G)$ the Frattini subgroup of G . If H is a subgroup of G , then $N_G(H)$ is the normalizer and $C_G(H)$ is the centralizer of H in G .

Lemma 2. *Let P be a metacyclic 2-group and $C_{2^k}^2 \cong Q < P$ with $k \geq 2$. Then the action of $N_P(Q)/Q$ on $Q/\Phi(Q)$ by conjugation is not faithful.*

Proof. By way of contradiction, we assume that $N_P(Q)/Q$ acts faithfully on $Q/\Phi(Q)$. Hence $|N_P(Q) : Q| = 2$, since $|\text{Aut}(Q/\Phi(Q))| = 6$. For simplicity, we may also assume $P = N_P(Q)$. Then $P/\Phi(Q)$ is a dihedral group of order 8. Let $\langle x \rangle \leq P$ such that $P/\langle x \rangle$ is cyclic. Then $\langle x \rangle \Phi(Q)/\Phi(Q)$ and $(P/\Phi(Q))/(\langle x \rangle \Phi(Q)/\Phi(Q)) \cong P/\langle x \rangle \Phi(Q)$ are also cyclic. Since a dihedral group of order 8 cannot have a cyclic quotient of higher order than 2, this shows $|P/\langle x \rangle \Phi(Q)| = 2$ and $|\langle x \rangle| = 4|\langle x \rangle \cap \Phi(Q)|$. Because $x^2 \in Q$, x has at most order 2^{k+1} . If x has lower order than 2^{k+1} , then x^2 would be contained in $\Phi(Q)$. Thus $|\langle x \rangle| = 2^{k+1}$. In particular $x \notin Q$. The restriction map $\varphi : \text{Aut}(\langle x \rangle) \rightarrow \text{Aut}(\langle x^2 \rangle)$ is an epimorphism with kernel of order 2. The action of Q on $\langle x \rangle$ induces a homomorphism $Q \rightarrow \text{Aut}(\langle x \rangle)$ with image contained in $\ker(\varphi)$, because $x^2 \in Q$. Since $\ker(\varphi)$ has order 2, x centralizes $\Phi(Q)$. By Burnside's basis theorem, there exists an element $y \in Q$ such that $Q = \langle y, xyx^{-1} \rangle$. Therefore $\Phi(Q) = \langle y^2, xy^2x^{-1} \rangle = \langle y^2 \rangle$ is cyclic. But this contradicts $k \geq 2$. □

3 Fusion systems

In this section, we will use the definitions and results of [4] to prove the main result of this paper. Moreover, we say that a fusion system $\mathcal{F}$ on a finite p -group P is *nilpotent* if $\text{Aut}_{\mathcal{F}}(Q)$ is a p -group for every $Q \leq P$ (cf. Theorem 3.11 in [4]).

Theorem 1. *Let P be a metacyclic 2-group, which is neither of maximal class nor homocyclic. Then every fusion system $\mathcal{F}$ on P is nilpotent.*

Proof. In order to show that P does not contain $\mathcal{F}$ -essential subgroups, let $Q \leq P$ be $\mathcal{F}$ -essential. Since $\text{Aut}_{\mathcal{F}}(Q)/\text{Aut}_Q(Q)$ has a strongly 2-embedded subgroup, $\text{Aut}(Q)$ is not a 2-group. By the Lemma 1, Q is homocyclic or a quaternion group of order 8. By Proposition 10.19 in [1], Q cannot be a quaternion group. Thus we may assume $Q \cong C_{2^k}^2$ for some $k \in \mathbb{N}$. Moreover, $C_P(Q) = Q$ holds, because Q is also $\mathcal{F}$ -centric. First, we consider the case $k = 1$. Then $Q \subseteq \langle x \in P : x^2 = 1 \rangle =: \Omega(P)$, and Exercise 1.85 in [1] imply $Q = \Omega(P) \leq P$. Since $P/Q = N_P(Q)/C_P(Q)$ is isomorphic to a subgroup of $\text{Aut}(Q)$, P has order 8. This contradicts our hypothesis. Hence, we have $k \geq 2$. Now consider the action of $\text{Aut}_{\mathcal{F}}(Q) = \text{Aut}_{\mathcal{F}}(Q)/\text{Aut}_Q(Q)$ on $Q/\Phi(Q)$. Lemma 2 shows that $1 \neq \text{Aut}_P(Q) \subseteq \text{Aut}_{\mathcal{F}}(Q)$ does not act faithfully. On the other hand, it is well-known that every automorphism of Q of odd order acts nontrivially on $Q/\Phi(Q)$. Therefore the kernel of the action under consideration forms a nontrivial normal 2-subgroup of $\text{Aut}_{\mathcal{F}}(Q)$, i.e. $O_2(\text{Aut}_{\mathcal{F}}(Q)) \neq 1$. But this contradicts the fact that $\text{Aut}_{\mathcal{F}}(Q)$ contains a strongly 2-embedded subgroup.

Now let Q be an arbitrary subgroup of P . We have to show that $\text{Aut}_{\mathcal{F}}(Q)$ is a 2-group. Let $\varphi \in \text{Aut}_{\mathcal{F}}(Q)$. Then Alperin's fusion theorem (Theorem 5.2 in [4]) shows that φ is the restriction of an automorphism of P . But again by Lemma 1, $\text{Aut}(P)$ is a 2-group, and φ must be a 2-element. This proves the theorem. □

Our last statement is in some sense a converse of Theorem 1.

Proposition 1. *Let P be a 2-group of maximal class or a homocyclic 2-group. Then there exists a fusion system on P , which is not nilpotent.*

Proof. It suffices to show that there exists a finite group G with P as a Sylow 2-subgroup such that G is not 2-nilpotent. If P is homocyclic, then the claim follows from Theorem 1.10 in [7]. If P has maximal class, then Theorem 2.4, Theorem 2.5, and Theorem 2.6 in [7] imply the result. $\square$

References

- [1] Y. Berkovich, *Groups of prime power order. Vol. 1*, de Gruyter Expositions in Mathematics, Walter de Gruyter GmbH & Co. KG, Berlin, 2008.
- [2] B. Huppert, *Endliche Gruppen I*, Die Grundlehren der Mathematischen Wissenschaften, Band 134, Springer-Verlag, Berlin, 1967.
- [3] S. Liedahl, *Enumeration of metacyclic p -groups*, J. Algebra **186** (1996), 436–446.
- [4] Markus Linckelmann, *Introduction to fusion systems*, in Group representation theory, 79–113, EPFL Press, Lausanne, 2007.
- [5] V. D. Mazurov, *Finite groups with metacyclic Sylow 2-subgroups*, Sibirsk. Mat. Ž. **8** (1967), 966–982.
- [6] Radu Stancu, *Control of fusion in fusion systems*, J. Algebra Appl. **5** (2006), 817–837.
- [7] R. W. Waall, *On p -nilpotent forcing groups*, Indag. Math. (N.S.) **2** (1991), 367–384.